

#### Ge Polyclonal Antibody to Homocysteine (HCy)

, Catalog # IC9984

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTICPROCEDURES

## [PROPERTIES]

Source: Polyclonal antibody preparation Host: Rabbit Purification: Antigen-specific Affinity Chromatography. Traits: Liquid Concentration: 200µg/mL UOM: 100µg Applications: WB; ICC; IHC-P; IHC-F; ELISA; IP; IF; FCM.

#### [IMMUNOGEN]

Immunogen: Small Molecule, HCy conjugated to BSA.

# [APPLICATIONS]

Western blotting: 0.5-2ug/ml Immunocytochemistry in formalin fixed cells: 5-20ug/ml Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml Optimal working dilutions must be determined by end user.

### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

Give us a call or send us a fax

info@immunoclone.com



Tel: 631-424-0089 Fax: 631-424-0095



Send a message to our company


# **I**C ImmunoClone

#### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles. Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



ImmunoClone(I&C) America P. O. Box 3988 Huntington Station, NY 55114 USA Give us a call or send us a fax

Tel: 631-424-**2**89 Fax: 631-424-0095



Send a message to our company

info@immunoclone.com

